



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 1 of 31

**Important Instructions to examiners:**

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more Importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answers and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 2 of 31

Q No.	Answer	marks
1	<b>Any 10</b>	<b>20</b>
1-a	<b>Characteristics of Organic compounds:</b> (any 2)  1. Generally found in living matter, i.e., animals and plants. 2. Insoluble in water, soluble in organic solvents. 3. Highly inflammable and volatile. 4. Poorer conductors of heat and electricity in aqueous solutions. 5. Usually always contain carbon, especially carbon-hydrogen bonds. 6. Classified into many classes on the basis of functional groups, known as homologous series. Each class is represented by a general formula and the members show similar properties 7. Examples: methane, ethane, acetylene, alcohols, carbon tetrachloride (CCl <sub>4</sub> ), urea	1 mark each
1-b	<b>IUPAC names:</b>  1. $\begin{array}{c} \text{OH} \\   \\ \text{CH}_3 - \text{CH} - \text{CH}_3 \end{array}$ 2- propanol  2. $\text{H}_3\text{C} - \text{CH}_2 - \text{CH} = \text{CH} - \text{CH}_2 - \text{CH}_2 - \text{CH}_3$ Heptene ( 3-Heptene)	1  1



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 3 of 31

1-c	<b>Physical properties of Alkane:(any 2)</b>  1) First four members(C1 to C4) of alkane are gases,next thirteen members (C5 to C17) are colourless liquids and higher alkanes are solids. 2) Liquid alkanes are lighter than water. 3) Alkanes are insoluble in water but freely soluble in organic solvent.  4) Boiling point and specific gravity increases with increase in molecular weight.	1 mark each						
1-d	<b>Uses of acetylene: (any 2)</b>  1. Ethyne is useful for artificial ripening and preservation of fruits. 2. It is useful in acetylene lamps to generate light. 3. Ethyne is used to prepare various organic compounds. 4. It is also useful to manufacture important organic compounds like acetic acid, acetaldehyde, ethyl alcohol and polymers like PVC...etc. 5. It is useful to produce an oxyacetylene flame. 6. is widely used as a fuel and a chemical building block. 7. Acetylene is used to volatilize carbon in radiocarbon dating. 8. Acetylene is sometimes used for carburization (that is, hardening) of steel when the object is too large to fit into a furnace	1 mark each						
1-e	<b>Products of the reaction:</b>  1. 1-3dibromopropane  2. Propane	1  1						
1-f	<b>Differences between alcohols &amp; phenol:</b> <table border="1"><thead><tr><th>PHENOL</th><th>ALCOHOL</th></tr></thead><tbody><tr><td>phenol reacts with FeCl<sub>3</sub> whereas normal</td><td>alcohol can not</td></tr><tr><td>Phenols are acidic</td><td>Alcohols are not acidic</td></tr></tbody></table>	PHENOL	ALCOHOL	phenol reacts with FeCl <sub>3</sub> whereas normal	alcohol can not	Phenols are acidic	Alcohols are not acidic	1 mark each for any 2
PHENOL	ALCOHOL							
phenol reacts with FeCl <sub>3</sub> whereas normal	alcohol can not							
Phenols are acidic	Alcohols are not acidic							



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 4 of 31

	Phenols are acidic and dissolve in a basic solution.	Alcohols are not acidic and will not dissolve in a basic solution	
	When phenol react with $\text{FeCl}_3$ it changes its colour from green to purple.	Alcohols produce no color change.	
	Phenols produce a brown tarry mass when combined with chromic acid	Not observe in alcohol	
	phenols dissolve in aqueous NaOH,	Not observe in alcohol	
	Aromatic	Aliphatic	
1-g	<b>Homologues of benzene:</b> Toluene( $\text{C}_6\text{H}_5\text{CH}_3$ ) Xylene( $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_3$ )		1 1
1-h	<b>Uses of phenol : (any two)</b> 1) The main use of phenol is as a feedstock for phenolic resins, bisphenol A and caprolactam (an intermediate in the production of nylon-6). 2) It is used in the manufacture of many products including insulation materials, adhesives, lacquers, paint, rubber, ink, dyes, illuminating gases, perfumes, soaps. 3) Also used in embalming and research laboratories. It is a product of the decomposition of organic materials, liquid manure, and the atmospheric degradation of benzene. 4) It is found in some commercial disinfectants, antiseptics, lotions and ointments. 5) Phenol is active against a wide range of microorganisms, and there are some medical and pharmaceutical applications including topical anaesthetic and ear drops, sclerosing agent. 6) It is used in dermatology for chemical face peeling.		1 mark each
1-i	<b>Nitration:</b> When benzene is treated with mixture of conc. $\text{H}_2\text{SO}_4$ and conc $\text{HNO}_3$ below		2



SUMMER-18 EXAMINATION  
Model Answer



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 6 of 31

	2) Liquid in liquid Example : Alcohol in water 3) Gas in liquid Example : water vapours in air, mist.							
1-1	<p><b>Indicator:</b> An indicator is an organic substance used in very small amount to determine the end point in a titration by a visual change of color.            Example:-methyl orange, phenolphthalein, starch, methyl red .</p> <p><b>Azeotropes:</b> are defined as the mixtures of liquids which boil at constant temperature like a pure liquid and possess same composition of components in liquid as well as in vapour phase. Azeotropes are also called constant boiling mixtures because whole of the azeotropes changes into vapour state at constant temperature and their components can not be separated by fractional distillation</p>	1            1						
<b>2</b>	<b>Any 4</b>	<b>16</b>						
2-a	<p><b>Homologous series-</b></p> <p>It is a series of class of organic compounds in which each member differs from its neighbor (immediate higher or lower member) by a constant difference of – CH<sub>2</sub>-(methylene) group.</p> <p>Homologous series of alkanes-CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, C<sub>3</sub>H<sub>8</sub>, C<sub>4</sub>H<sub>10</sub>, C<sub>5</sub>H<sub>12</sub></p> <p>From above example propane(C<sub>3</sub>H<sub>8</sub>) differs from ethane (C<sub>2</sub>H<sub>6</sub>) or butane(C<sub>4</sub>H<sub>10</sub>) by a –CH<sub>2</sub>- group.</p> <p>Every class of organic compounds has such a homologous series. All members of homologous series have the same general formula. Each homologous series has a particular functional group.</p>	4						
2-b	<p><b>Classification organic compounds on the basis of functional group:</b></p> <table border="1" style="width:100%; margin-top:10px;"> <thead> <tr> <th style="width:30%;">Functional Group</th> <th style="width:30%;">Name</th> <th style="width:40%;">Example</th> </tr> </thead> <tbody> <tr> <td style="height:30px;"></td> <td></td> <td></td> </tr> </tbody> </table>	Functional Group	Name	Example				4
Functional Group	Name	Example						



SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 7 of 31

	Alkane	CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub> (propane)
	Alkene	CH <sub>3</sub> CH=CH <sub>2</sub> (propene)
	Alkyne	CH <sub>3</sub> C≡CH (propyne)
F, Cl, Br, or I	Alkyl halide	CH <sub>3</sub> Br (methyl bromide)
	Alcohol	CH <sub>3</sub> CH <sub>2</sub> OH (ethanol)
	Ether	CH <sub>3</sub> OCH <sub>3</sub> (dimethyl ether)
	Amine	CH <sub>3</sub> NH <sub>2</sub> (methyl amine)
	Aldehyde	CH <sub>3</sub> CHO (acetaldehyde)
	Ketone	CH <sub>3</sub> COCH <sub>3</sub> (acetone)
	Acyl chloride	CH <sub>3</sub> COCl (acetyl chloride)
	Carboxylic acid	CH <sub>3</sub> CO <sub>2</sub> H (acetic acid)
	Ester	CH <sub>3</sub> CO <sub>2</sub> CH <sub>3</sub> (methyl acetate)







SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 8 of 31

	<table border="1"><tr><td></td><td>Amide</td><td>CH<sub>3</sub>NH<sub>2</sub> (acetamide)</td></tr></table>		Amide	CH <sub>3</sub> NH <sub>2</sub> (acetamide)	
	Amide	CH <sub>3</sub> NH <sub>2</sub> (acetamide)			
2-c	<p><b>Aromatisation in alkanes:</b> The process of conversion of aliphatic compound into aromatic compound is known as <b>aromatization</b>. <b>Alkanes</b> having six to 10 carbon atoms are converted into benzene and its homologues at high pressure and temperature in presence of catalyst.</p> <p>Ex.</p> 	4			
2-d	<p><b>Methods of preparation of alkenes:</b> (Any 2)</p> <p>1) <b>By Dehydration of alcohols :</b></p> <p>When alcohols methods is heated in presence of sulphuric acide , water is eliminated and alkene is formed.</p> $\begin{array}{c} \text{H}_3\text{C}-\text{CH}-\text{CH}_2 \\   \quad   \\ \text{H} \quad \text{OH} \\ \text{- Propanol} \end{array} \xrightarrow[170^\circ\text{c}]{\text{H}_2\text{SO}_4} \text{CH}_3-\text{CH}=\text{CH}_2 + \text{H}_2\text{O}$ <p style="text-align: center;">propene</p> <p>2) <b>By dehydrogenation of alkyl halide :</b></p> <p>When alkyl halide heated with alcoholic solution of Na or k-hydroxide, hydrogen halide is eliminated and alkene is formed.</p>	2 marks each for any 2			






SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry


Subject code : 17312

Page 9 of 31





3) **By dehalogenation of vicinal dihalide. :**

A compound having two halogen atoms on adjacent carbon atoms is called a vicinal dihalide. Alkenes are formed when vicinal dihalides are heated with Zn-dust in ethyl alcohol.



4) **By cracking of Alkanes:**

Alkanes when heated at 500 – 700 °c in absence of air, decomposes to yield lower molecular weight alkenes, alkanes and hydrogen.



2-e

**Isomerism in alkyl halides:** The isomerism exhibited by alkyl halides is of two types: Chain isomerism and Position isomerism

**Chain isomerism:** Paraffins exhibit chain isomerism. Alkyl halides being derivatives of paraffins show chain isomerism depending upon the nature of the

4



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 10 of 31

	<p>chain whether it is straight or branched. For example, difference between the following bromo derivatives of butane is due to difference in the nature of the chain.</p> <p><math>\text{CH}_3 \cdot \text{CH}_2 \cdot \text{CH}_2 \cdot \text{CH}_2\text{Br}</math> n butyl bromide</p> <p><math>\text{CH}_3 - \text{CH} - \text{CH}_2\text{Br}</math> Isobutyl bromide   <math>\text{CH}_3</math></p> <p><b>Position isomerism:</b> Another type of isomerism exhibited by alkyl halide is due to the difference in the position of the halogen atom in the same chain and is termed as position isomerism.</p> <p>Ex. <math>\text{CH}_3 \cdot \text{CH}_2 \cdot \text{CH}_2\text{I}</math> n- propyl iodide</p> <p>And <math>\text{CH}_3 \cdot \text{CH I} \cdot \text{CH}_3</math> Isopropyl iodide</p> <p>Both have straight chain formula but differ in the position of the halogen atom.</p>	
2-f	<p><b>Differentiate between primary, secondary and tertiary alcohols.</b></p> <p>To differentiate between primary, secondary and tertiary alcohols, four methods are used</p> <p>(1) Oxidation method (2) Action of hot reduced cu. (3) victor mayer's method (4) Lucas test.</p> <p>1) <b>Oxidation method</b> :- i) primary alcohols easily oxidized to aldehydes and then to acids, containing same number of carbon atoms , as the original alcohol.</p>	2 marks each for any 2




SUMMER-18 EXAMINATION  
Model Answer


Subject title: Industrial Chemistry

Subject code : 17312


Page 11 of 31




ii) Secondary alcohol, gives ketone, which on prolonged oxidation, by action of oxidizing agent gives acids.



iii) Tertiary alcohol are oxidized by acid oxidizing agents to give mixture of ketone and acid



(2) By action of hot reduced Cu : with hot reduced Cu at 300°C, primary alcohol gives aldehydes secondary gives ketone and tertiary alcohols gives olefins.





SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 12 of 31

	<p>Isopropyl alcohol (secondary)</p> $\begin{array}{c} \text{CH}_3 \\   \\ \text{CH}_3-\text{CHOH} \\   \\ \text{CH}_3 \end{array} \xrightarrow{\text{Cu}/300^\circ\text{C}} \begin{array}{c} \text{CH}_3-\text{C}=\text{CH}_2 \\   \\ \text{CH}_3 \end{array} + \text{H}_2\text{O}$ <p>Tertiary butyl alcohol isobutylene</p> <p><i>Note: Other two methods are also to be considered</i></p>	
3	<b>Any 4</b>	<b>16</b>
3-a	<p>(i) <b>2,3 dimethyl heptane</b> <math>\text{H}_3\text{C}-\underset{\text{I}}{\text{CH}}-\underset{\text{I}}{\text{CH}}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_3</math> <math>\text{CH}_3 \quad \text{CH}_3</math></p> <p>(ii) <b>2-methyl propane</b> <math>\text{H}_3\text{C}-\underset{\text{I}}{\text{CH}}-\text{CH}_3</math> <math>\text{CH}_3</math></p>	<p>2</p> <p>2</p>
3-b	<p><b>Bayers strain theory-</b> This theory based on the fact that the normal angle between any pair of bonds of carbon atom is <math>109^\circ 28'</math>. Baeyer postulated that any deviation of bond angles from the normal tetrahedral value impose condition of internal strain on the ring. He also assumed that all cycloalkanes were planar &amp; calculated the angles through which each of the valency bond was deflected from the normal direction in the formation of various rings. It is called angle strain which</p>	4







SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 13 of 31

determine stability of ring.


1.	Cyclopropane		60°
2.	Cyclobutane		90°
3.	Cyclopentane		108°
4.	Cyclohexane		120°

In cyclopropane three c atoms occupy the corners of an equilateral triangle. Therefore bond angles of 60°. Normal angle 109°28' is compressed to 60° & that each of the two bonds involved is pulled in by

$$\frac{1}{2}(109^{\circ}28' - 60^{\circ})$$

$$\frac{1}{2}(49^{\circ}28')$$

$$=24^{\circ}44'$$



Angle strain in cyclopropane

The value represents angle strain or the deviation through which each bond



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 14 of 31

bends from the normal value.


1.	Cyclopropane		24° 44'
2.	Cyclobutane		9° 44'
3.	Cyclopentane		0° 44'
4.	Cyclohexane		-5° 16'

The angle strain for other cycloalkanes can be calculated in the same way.

The angle strain is maximum in case of cyclopropane. According to this theory cyclopropane should be highly strained molecule & unstable also.

3-c

**Structure of benzene-**



**Preparation method of benzene-**

1) By heating an aromatic acid or its sodium salt with sodalime (Removal of COOH group)

2

1 mark  
each for  
any 2  
methods



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 15 of 31

	$C_6H_5COONa + NaOH \longrightarrow C_6H_6 + Na_2CO_3$ <p>Sodium benzoate    benzene</p> <p>2)By the hydrolysis of sulphonic acid with superheated steam</p> $C_6H_5SO_3H + H_2O \longrightarrow C_6H_6 + H_2SO_4$ <p>Benzene sulphonic    benzene acid</p> <p><i>Note: Any other method should be given mark</i></p>	
3-d	<p><b>Action of ammonia on phenol</b></p> <p>When phenol is heated with ammonia in presence of anhydrous zinc chloride then formation of Aniline takes place</p> $  \begin{array}{c} \text{OH} \\   \\ \text{C}_6\text{H}_5 \end{array} + \text{NH}_3 \xrightarrow{\text{ZnCl}_2} \begin{array}{c} \text{NH}_2 \\   \\ \text{C}_6\text{H}_5 \end{array} + \text{H}_2\text{O}  $ <p><b>Action of H<sub>2</sub>SO<sub>4</sub> on phenol –</b></p> <p>Phenol when treated with conc. H<sub>2</sub>SO<sub>4</sub> at room temp. gives o-phenol sulphonic acid.</p> $  \begin{array}{c} \text{OH} \\   \\ \text{C}_6\text{H}_5 \end{array} + \text{HO-SO}_3\text{H} \xrightarrow{15-20^\circ\text{C}} \begin{array}{c} \text{OH} \\   \\ \text{C}_6\text{H}_4 \\   \\ \text{SO}_3\text{H} \end{array} + \text{H}_2\text{O}  $ <p>Phenol    sulfuric acid    o-phenol sulphonic acid</p>	2
3-e	<p><b>Chain isomerism</b>-Alcohol show chain isomerism depending upon nature of chain i.e. straight or branched.</p> <p>CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>OH n-butyl alcohol</p> <p>CH<sub>3</sub>-CH-CH<sub>2</sub>-OH</p> <p style="margin-left: 40px;">I</p> <p style="margin-left: 40px;">CH<sub>3</sub></p>	2



SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 16 of 31

	<p>The isomerism is due to the difference in the nature of chain.i.e.straight or branched</p> <p><b>Position isomerism</b>-It is due to the difference in the position of hydroxyl group in the same chain .</p> <p>Both have straight chain formula but differ in the position of hydroxyl group.</p> <p><b>Functional isomerism</b>-Monohydric alcohols are isomeric with ethers ,general formula for both of these <math>C_nH_{2n+2}O</math>.So ethanol <math>C_2H_5OH</math> is isomeric with dimethyl ether,<math>CH_3.O.CH_3</math> &amp; propanol,<math>C_3H_7OH</math> is isomeric with ethylmethyl ether <math>C_2H_5.O.CH_3</math>.This type of isomerism is due to the presence of different functional groups.</p>	1          1
3-f	<p><b>Quinonoid theory:</b> This theory explains the color changes in titrations on the basis of intramolecular changes.</p> <p>This theory believes that-</p> <ol style="list-style-type: none"><li>1. An acid base indicator is either a weak acid or a weak base.</li><li>2. An indicator consists of an equilibrium mixture of at least two tautomeric forms one is benzenoid while the other is Quinonoid form.</li><li>3. The two forms posses different colors.</li><li>4. Out of these forms, one exists in an acid solution &amp; the other form exists in an alkaline solution.</li><li>5. The Quinonoid form is generally deeper in colour than beenzenoid form.</li><li>6.As the PH of the solution containing an indicator changes one form of the indicator changes to the other &amp; as a result of this the solution shows a change of colour</li></ol> <p>There are two tautomeric forms of methyl orange .The Quinonoid form (red form)exists in an acidic solution &amp; it passes to the Benzenoid form (yellow) as the PH changes to the alkaline side.</p> <p>Phenolphthalein is colorless in an acidic solution where it exists in the</p>	4





SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 17 of 31

	<p>Benzenoid form .If an alkali is added, it changes to theQuinonoid form &amp; imparts pink color to the solution</p> <p>Benzenoid form                      Quinonoid form</p>	
4	<b>Any 4</b>	<b>16</b>
4-a	<p>Organic compounds are classified into four categories on the basis of structure.</p> <ol style="list-style-type: none"><li><b>Aliphatic compounds :</b> Which consist of open chain of carbon atoms are called aliphatic compounds. Example : (Any 1) a) Propane <math>\text{CH}_3\text{-CH}_2\text{-CH}_3</math> b) Acetic acid <math>\text{CH}_3\text{-COOH}</math></li><li><b>Alicyclic compounds:</b> These are cyclic compounds composed of ring of carbon atoms with properties similar to aliphatic compounds. Example : (Any 1)</li></ol> <p>a)  OR </p> <p>cyclopropane</p>	4




SUMMER-18 EXAMINATION  
Model Answer


Subject title: Industrial Chemistry

Subject code : 17312


Page 18 of 31




OR



Cyclobutane



OR




Cyclohexane

3. Aromatic compounds:


These are cyclic compounds having six membered ring of carbon atoms with alternate single and double bonds

Example :

a) Benzene



b) Toluene





SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312


Page 19 of 31

4. Heterocyclic compounds:


These are cyclic compounds in which ring consist of atoms atoms and some other element such as oxygen, nitrogen, sulphur.

Example : (Any 1)

a) pyridine



b) pyrrole



c) furan



d) Thiopheno



4-b

**Saturated compounds** –Hydrocarbons in which all the bonds of carbon atoms are fully utilized & cannot take up more hydrogen atoms are known as saturated compounds.

e.g. alkanes  $\text{CH}_4$ -methane

**Unsaturated hydrocarbons** – Hydrocarbons in which all the bonds of carbon atoms are not fully utilized & can take up more hydrogen atoms are known as unsaturated hydrocarbons. The carbon atoms are join to each other by double or triple bond.

e.g. Alkenes & Alkynes( $\text{C}_2\text{H}_4$ ,  $\text{C}_2\text{H}_2$ )

1

1

1

1







SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 21 of 31

	<p>Examples of ideal solutions-Hexane + Heptane,Benzene + Toluene</p> <p>2. Non ideal solution –The solutions which do not obey Raoult's law. Like forces are not similar to unlike forces.Change in volume &amp; enthalpy on mixing the two liquid.There are two types of non ideal solutions-solutions which show positive deviation from Raoult's law &amp; negative deviation from Raoult's law.</p> <p>Examples-positive deviation-Acetone + carbon disulphide Negative deviation –Chloroform + acetone</p>	
4-f	<p><b>Minimum boiling azeotropes</b>-is solution of definite compositions which boils at a definite temp.which is lower than the B.P.of both the components of solution.</p> <p>Minimum boiling azeotropes show positive deviation from Raoult's law.</p> <p>Examples of minimum boiling azeotrope /azeotropic mixture-ethanolwater, chloroform-methanol etc.</p> <div style="display: flex; justify-content: space-around; align-items: center;"><div style="text-align: center;"><p>Boiling point diagram</p></div><div style="text-align: center;"><p>equilibrium diagram</p></div></div> <p><b>Maximum boiling azeotrope</b> is a solution of definite composition which boils at a temp. higher than the B.P. of both.the components of the solution</p>	2




**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 22 of 31

	<p>Maximum boiling azeotropes show negative deviation from Raoult's law.</p> <p>Examples of maximum boiling azeotrope-HCl-water, water-nitric acid</p>  <p>Boiling point diagram                      equilibrium diagram</p>	2
5	<b>Any 4</b>	<b>16</b>
5-a	<p><b>IUPAC Rules for naming monofunctional compounds:</b></p> <ol style="list-style-type: none"><li>1) Identify the principal functional group.</li><li>2) Number the longest chain containing the functional group from the end closer to it.</li><li>3) Write the parent name corresponding to the number of carbons in the longest chain.</li><li>4) arrange the substituent names with position numbers in alphabetic order</li><li>5) Prefix substituent's name with the parent name.</li></ol>	4
5-b	<p><b>Phenol from Cumene:</b></p> <p>By oxidation of Cumene followed by treatment with acid we can produce phenol. It is generally used for manufacture of phenol.</p>	2



SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 23 of 31

	<p style="text-align: center;">Cumene      Cumene hydroperoxide      phenol</p>	2											
5-c	<p><b>Physical properties of alcohol:</b></p> <ol style="list-style-type: none"> <li>1) Higher members are colorless waxy solids.</li> <li>2) Lower members are soluble in water &amp; organic solvents.</li> <li>3) These are lighter than water.</li> <li>4) Lower members have pleasant smell while higher member are odorless &amp; tasteless.</li> </ol>	1 mark each											
5-d	<p><b>Distinguish between ideal and non ideal solution:</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th data-bbox="289 1360 654 1528" rowspan="2">Ideal solutions</th> <th colspan="2" data-bbox="654 1360 1317 1415">Non-ideal solutions</th> </tr> <tr> <th data-bbox="654 1415 964 1528">Positive deviation from Raoult's law</th> <th data-bbox="964 1415 1317 1528">Negative deviation from Raoult's law</th> </tr> </thead> <tbody> <tr> <td data-bbox="289 1528 654 1696">1. Obey Raoult's law at every range of concentration.</td> <td data-bbox="654 1528 964 1696">1. Do not obey Raoult's law.</td> <td data-bbox="964 1528 1317 1696">1. Do not obey Raoult's law.</td> </tr> <tr> <td data-bbox="289 1696 654 1864">2. <math>\Delta H_{mix} = 0</math>; neither heat is evolved nor absorbed during</td> <td data-bbox="654 1696 964 1864">2. <math>\Delta H_{mix} &gt; 0</math>. Endothermic dissolution; heat is</td> <td data-bbox="964 1696 1317 1864">2. <math>\Delta H_{mix} &lt; 0</math>. Exothermic dissolution; heat is evolved.</td> </tr> </tbody> </table>	Ideal solutions	Non-ideal solutions		Positive deviation from Raoult's law	Negative deviation from Raoult's law	1. Obey Raoult's law at every range of concentration.	1. Do not obey Raoult's law.	1. Do not obey Raoult's law.	2. $\Delta H_{mix} = 0$ ; neither heat is evolved nor absorbed during	2. $\Delta H_{mix} > 0$ . Endothermic dissolution; heat is	2. $\Delta H_{mix} < 0$ . Exothermic dissolution; heat is evolved.	1 mark each for any 4
Ideal solutions	Non-ideal solutions												
	Positive deviation from Raoult's law	Negative deviation from Raoult's law											
1. Obey Raoult's law at every range of concentration.	1. Do not obey Raoult's law.	1. Do not obey Raoult's law.											
2. $\Delta H_{mix} = 0$ ; neither heat is evolved nor absorbed during	2. $\Delta H_{mix} > 0$ . Endothermic dissolution; heat is	2. $\Delta H_{mix} < 0$ . Exothermic dissolution; heat is evolved.											



SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 24 of 31

dissolution.	absorbed.	
3. $\Delta V_{mix} = 0$ ; total volume of solution is equal to sum of volumes of the components.	3. $\Delta V_{mix} > 0$ . Volume is increased after dissolution.	3. $\Delta V_{mix} < 0$ . Volume is decreased during dissolution.
4. $P = p_A + p_B = p_A^0 X_A + p_B^0 X_B$ i.e., $p_A = p_A^0 X_A ; p_B = p_B^0 X_B$	4. $p_A > p_A^0 X_A ;$ $p_B > p_B^0 X_B \therefore$ $p_A + p_B > p_A^0 X_A + p_B^0 X_B$	4. $p_A < p_A^0 X_A ; p_B < p_B^0 X_B$ $\therefore$ $p_A + p_B < p_A^0 X_A + p_B^0 X_B$
5. $A-A, A-B, B-B$ Interactions should be same, i.e., 'A' and 'B' are identical in shape, size and character.	5. $A-B$ Attractive force should be weaker than $A-A$ and $B-B$ attractive forces. 'A' and 'B' have different shape, size and character.	5. $A-B$ Attractive force should be greater than $A-A$ and $B-B$ attractive forces. 'A' and 'B' have different shape, size and character.
6. Escaping tendency of 'A' and 'B' should be same in pure liquids and in the solution.	6. 'A' and 'B' escape easily showing higher vapour pressure than the expected value.	6. Escaping tendency of both components 'A' and 'B' is lowered showing lower vapour pressure than expected ideally.
7.Examples: benzene + toluene; n-hexane + n-heptane;	7. Examples: Acetone + ethanol	7. Examples: Acetone + aniline;





SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 25 of 31

5-e	<b>Methods of choosing indicators for acid- alkali titration:</b>	4									
	<ul style="list-style-type: none"><li>• an appropriate indicator will change colour at the equivalence point of the titration.  Litmus is not used in titrations because the pH range over which it changes colour is too great.</li><li>• Universal indicator which is actually a mixture of several indicators displays a variety of colours over a wide pH range so it can be used to determine an approximate pH of a solution but is not used for titrations. an appropriate indicator will change colour at the equivalence point of the titration.  Litmus is not used in titrations because the pH range over which it changes colour is too great.</li><li>• Determine what species are present at the equivalence point &amp; deduce the pH at the equivalence point</li></ul>										
	<table border="1"><thead><tr><th data-bbox="355 1478 813 1675"><b>pH of salts formed from reactions of acids &amp; bases (25°C)</b></th><th data-bbox="818 1478 1040 1675"><b>Strong Base</b></th><th data-bbox="1045 1478 1247 1675"><b>Weak Base</b></th></tr></thead><tbody><tr><td data-bbox="355 1675 813 1770"><b>Strong Acid</b></td><td data-bbox="818 1675 1040 1770">pH = 7</td><td data-bbox="1045 1675 1247 1770">pH &lt; 7</td></tr><tr><td data-bbox="355 1770 813 1852"><b>Weak Acid</b></td><td data-bbox="818 1770 1040 1852">pH &gt; 7</td><td data-bbox="1045 1770 1247 1852">pH = 7</td></tr></tbody></table>	<b>pH of salts formed from reactions of acids &amp; bases (25°C)</b>	<b>Strong Base</b>	<b>Weak Base</b>	<b>Strong Acid</b>	pH = 7	pH < 7	<b>Weak Acid</b>	pH > 7	pH = 7	
<b>pH of salts formed from reactions of acids &amp; bases (25°C)</b>	<b>Strong Base</b>	<b>Weak Base</b>									
<b>Strong Acid</b>	pH = 7	pH < 7									
<b>Weak Acid</b>	pH > 7	pH = 7									



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry


Subject code : 17312

Page 26 of 31

- Use the table of indicators to choose an indicator which changes colour over a pH range that includes the equivalence point

***Examples:***

**Strong Acid - Strong Base titration**




At equivalence the only species present will be NaCl(aq) & H<sub>2</sub>O(l)

The solution of a salt of a strong acid and a strong base will have a pH=7

NaCl(aq) will have a pH=7

A suitable indicator would be bromothymol blue (pH range 6.2 - 7.6) or phenol red (pH range 6.8 - 8.4)

**Strong Acid - Weak Base titration**




NH<sub>4</sub>Cl is the salt of a strong acid & a weak base, so a solution of NH<sub>4</sub>Cl will have a pH < 7 (NH<sub>4</sub><sup>+</sup> is a weak acid)

A suitable indicator would be methyl orange (pH range 3.1 - 4.4) or methyl red

(pH range 4.4 - 6.0)

**Weak Acid - Strong Base titration**



CH<sub>3</sub>COONa is the salt of a weak acid & a strong base, so a solution of CH<sub>3</sub>COONa will have a pH > 7 (CH<sub>3</sub>COO<sup>-</sup> is a weak base)

A suitable indicator would be phenolphthalein (pH range 8.3 - 10.0) or





SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 27 of 31

	thymol blue (pH 8.0 - 9.6)	
5-f	<p><b>Methods for preparation of cycloalkanes (any one)</b></p> <p><b>i) From Di-halides :</b></p> <p>Terminal di-halide are treated with sodium or zinc to form cycloalkanes.</p>  <p><b>ii) From aromatic hydrocarbons :</b></p>  <p><b>physical properties of cycloalkanes:</b></p> <ol style="list-style-type: none"><li>1. Cyclopropane and cyclobutane are gases at room temperature. the remaining cycloalkanes are liquids.</li><li>2. melting and boiling points of cycloalkane increases with increase of</li></ol>	2



SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 28 of 31

	molecular weight. 3. Cycloalkanes are insoluble in water but dissolves in ethanol and water.	
<b>6</b>	<b>Any four</b>	<b>16</b>
6-a	<b>Polymerisation :</b> It is the process of combination of two or more monomeric units to form a high molecular weight compound with or without the elimination of H <sub>2</sub> O , HCl etc. under specific conditions of temperature pressure and catalystr. Example : a) $n\text{CH}_2=\text{CH}_2$ ethylene	2  2
6-b	<b>Friedel-crafts reaction:</b> Benzene reacts at room temperature with a chloro alkane (for example, chloromethane or chloro ethane) in the presence of aluminium chloride as a catalystr. It is known as <b>Friedel-Crafts rection of benzene</b>  Benzene  Toluene  <b>Grignard's reaction:</b>	2




SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 29 of 31

		2
6-c	<p><b>Methods of preparation of alkenes:</b></p> <p><b>By Dehydration of alcohols :</b></p> <p>When alcohols are heated in the presence of sulphuric acid, water is eliminated and alkene is formed.</p> $\begin{array}{ccc} \text{H}_3\text{C}-\text{CH}-\text{CH}_2 & \xrightarrow[170^\circ\text{C}]{\text{H}_2\text{SO}_4} & \text{CH}_3-\text{CH}=\text{CH}_2 + \text{H}_2\text{O} \\   \quad   & & \\ \text{H} \quad \text{OH} & & \\ \text{- Propanol} & & \text{propene} \end{array}$ <p><b>By dehydrogenation of alkyl halide :</b></p> <p>When alkyl halide is heated with alcoholic solution of Na or K-hydroxide, hydrogen halide is eliminated and alkene is formed.</p> $\begin{array}{ccc} \text{H}_3\text{C}-\text{CH}-\text{CH}_2 + \text{KOH} & \xrightarrow[\text{Alcohol}]{\Delta} & \text{CH}_3-\text{CH}=\text{CH}_2 + \text{KBr} + \text{H}_2\text{O} \\   \quad   & & \\ \text{H} \quad \text{Br} & & \\ \text{- Bromopropane} & & \text{propene} \end{array}$	2
6-d	<p><b>Ostwald theory :</b> Ostwald's theory considers indicator to be a weak acid or base whose unionised forms differently coloured. In presence of acid or base, i.e. pH change, there is ionization of indicator and hence the colour change appears.</p> <p>For example <a href="#">phenolphthalein</a></p>	4



**SUMMER-18 EXAMINATION**  
**Model Answer**

Subject title: Industrial Chemistry

Subject code : 17312

Page 30 of 31

	<p>phenolphthalein is a weak acid (PhH)</p> $\text{PhH} \rightleftharpoons \text{Ph}^- + \text{H}^+ \dots\dots\dots(1)$ <p>(colourless (Pink in base) in acid)</p> $\text{H}^+ + \text{OH}^- \rightleftharpoons \text{H}_2\text{O}$ <p>In presence of an acid (H<sup>+</sup>) equilibrium (1) is displaced towards the left hand side (a case of LeChatelier's principle); when strong base like NaOH is added, this equilibrium is displaced towards right hand side and there is colour change from colourless to pink when pH changes. This indicator is not suitable for titrating weak base since weak base can't furnish enough OH<sup>-</sup> that can react with H<sup>+</sup> of the phenolphthalein and can impart pink colour only after excess of weak base is added.</p>	
6-e	<p><b>Azeotropic Mixture</b></p> <p>Azeotropes are defined as the mixtures of liquids which boil at constant temperature like a pure liquid and possess same composition of components in liquid as well as in vapour phase. Their dew point and bubble point are identical. Azeotropes are also called constant boiling mixtures because whole of the azeotropes changes into vapour state at constant temperature and their components can not be separated by fractional distillation.</p> <p>Azeotrope can not be separated by distillation because the dew point and bubble point are identical.</p>	4



SUMMER-18 EXAMINATION  
Model Answer

Subject title: Industrial Chemistry

Subject code : 17312

Page 31 of 31

6-f	<p><b>Ethylamine from ethyl iodide:</b></p> <p>When alkyl halides are heated with ethanoic solution of ammonia under pressure in a sealed tube a mixture of amines are obtained.</p> $\text{C}_2\text{H}_5\text{I} + \text{H NH}_2 \longrightarrow \text{C}_2\text{H}_5\text{NH}_2 + \text{HI}$	4
-----	---	---