



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22206

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answer and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

Q. No.	Sub Q. N.	Answer	Marking Scheme
1.		Attempt any FIVE of the following:	10
	a)	State whether the function $f(x) = \frac{a^x + a^{-x}}{2}$ is even or odd .	02
	Ans	$f(x) = \frac{a^x + a^{-x}}{2}$ $\therefore f(-x) = \frac{a^{-x} + a^{-(-x)}}{2}$ $= \frac{a^{-x} + a^x}{2}$ $= f(x)$ $\therefore \text{function is even.}$	1 ½ ½
b)	If $f(x) = x^2 + 6x + 10$ find $f(2) + f(-2)$	02	
Ans	$f(x) = x^2 + 6x + 10$ $\therefore f(2) = (2)^2 + 6(2) + 10 = 26$ $\therefore f(-2) = (-2)^2 + 6(-2) + 10 = 2$ $\therefore f(2) + f(-2) = 26 + 2 = 28$	½ ½ 1	
c)	If $y = \log(x^2 + 2x + 5)$, find $\frac{dy}{dx}$	02	
Ans	$y = \log(x^2 + 2x + 5)$ $\therefore \frac{dy}{dx} = \frac{1}{x^2 + 2x + 5} (2x + 2)$	02	



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
1.		$\therefore \frac{dy}{dx} = \frac{2x+2}{x^2+2x+5}$	
	d)	Evaluate : $\int \frac{1}{\sin^2 x \cos^2 x} dx$	02
	Ans	$\int \frac{1}{\sin^2 x \cos^2 x} dx$ $= \int \frac{\sin^2 x + \cos^2 x}{\sin^2 x \cos^2 x} dx$ $= \int \frac{1}{\cos^2 x} + \frac{1}{\sin^2 x} dx$ $= \int (\sec^2 x + \operatorname{cosec}^2 x) dx$ $= \tan x - \cot x + c$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
		OR	
		$\int \frac{1}{\sin^2 x \cos^2 x} dx$ $= \int \operatorname{cosec}^2 x \cdot \sec^2 x dx$ $= \int (1 + \cot^2 x)(1 + \tan^2 x) dx$ $= \int (1 + \tan^2 x + \cot^2 x + \tan^2 x \cot^2 x) dx$ $= \int (1 + \tan^2 x + \cot^2 x + 1) dx$ $= \int (\sec^2 x + \operatorname{cosec}^2 x) dx$ $= \tan x - \cot x + c$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
	e)	Find the area enclosed by the curve $y = 3x^2$, x -axis and the ordinates $x = 1$, $x = 3$	02
	Ans	$\text{Area } A = \int_a^b y dx$ $= \int_1^3 3x^2 dx$ $= 3 \left[\frac{x^3}{3} \right]_1^3 \quad \text{OR} = \left[x^3 \right]_1^3$ $= 3 \left[\frac{3^3}{3} - \frac{1^3}{3} \right] = \left[3^3 - 1^3 \right]$ $= 26$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22206

Q. No.	Sub Q. N.	Answer	Marking Scheme
1.	f)	An unbiased coin is tossed 5 times .Find the probability of getting a head.	02
	Ans	$n = 5, p = \frac{1}{2}, q = \frac{1}{2}, r = 1$ $p(r) = {}^n C_r (p)^r (q)^{n-r}$ $p(1) = 5 {}^5 C_1 \left(\frac{1}{2}\right)^1 \left(\frac{1}{2}\right)^{5-1}$ $= \frac{5}{32} \text{ or } 0.156$	<p>½</p> <p>½</p> <p>1</p>
	g)	Evaluate: $\int x \cos x dx$	02
	Ans	$\int x \cos x dx = x \int \cos x dx - \int \left(\int \cos x dx \cdot \frac{d}{dx} x \right) dx$ $= x \sin x - \int (\sin x \cdot 1) dx$ $= x \sin x + \cos x + c$	<p>½</p> <p>1</p> <p>½</p>
2		Attempt any THREE of the following:	12
	(a)	If $e^x + e^y = e^{x+y}$, find $\frac{dy}{dx}$	04
	Ans	$e^x + e^y = e^{x+y}$ $e^x + e^y \frac{dy}{dx} = e^{x+y} \left[1 + \frac{dy}{dx} \right]$ $e^y \frac{dy}{dx} - e^{x+y} \frac{dy}{dx} = e^{x+y} - e^x$ $\frac{dy}{dx} (e^y - e^{x+y}) = e^{x+y} - e^x$ $\frac{dy}{dx} = \frac{e^{x+y} - e^x}{e^y - e^{x+y}}$	<p>1</p> <p>1</p> <p>1</p> <p>1</p>
	b)	If $x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{2}$	04
	Ans	$x = a(\theta + \sin \theta)$ $\frac{dx}{d\theta} = a(1 + \cos \theta)$ $y = a(1 - \cos \theta)$	1



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme	
2.	b)	$\frac{dy}{d\theta} = a(-(-\sin \theta)) = a \sin \theta$	1	
		$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{a \sin \theta}{a(1 + \cos \theta)}$		
		$= \frac{\sin \theta}{(1 + \cos \theta)}$ OR $= \frac{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}} = \tan \frac{\theta}{2}$	½	
		at $\theta = \frac{\pi}{2}$, $\frac{dy}{dx} = \frac{\sin \frac{\pi}{2}}{(1 + \cos \frac{\pi}{2})} = \tan \frac{\pi}{4}$	½	
			$= \frac{1}{1+0} = 1$ =1	1

	c)	Find the maximum and minimum values of $y = 2x^3 - 3x^2 - 36x + 10$	04	
	Ans	Let $y = 2x^3 - 3x^2 - 36x + 10$		
		$\therefore \frac{dy}{dx} = 6x^2 - 6x - 36$	½	
		$\therefore \frac{d^2y}{dx^2} = 12x - 6$	½	
	Consider $\frac{dy}{dx} = 0$			
	$6x^2 - 6x - 36 = 0$			
	$x^2 - x - 6 = 0$			
	$\therefore x = -2, x = 3$	1		
	at $x = -2$			
	$\frac{d^2y}{dx^2} = 12(-2) - 6 = -30 < 0$	½		
	$\therefore y$ is maximum at $x = -2$			
	$y_{\max} = 2(-2)^3 - 3(-2)^2 - 36(-2) + 10$			
	$= 54$	½		
	at $x = 3$, $\frac{d^2y}{dx^2} = 12(3) - 6 = 30 > 0$	½		
	$\therefore y$ is minimum at $x = 3$			
	$y_{\min} = 2(3)^3 - 3(3)^2 - 36(3) + 10$			
	$= -71$	½		



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
3.	Ans	$y = 2x - x^2$	
		$\frac{dy}{dx} = 2 - 2x$	
		at (2,0)	
		slope of tangent $m = \frac{dy}{dx} = 2 - 2(2) = -2$	
		equation of tangent is,	
		$y - y_1 = m(x - x_1)$	
		$y - 0 = -2(x - 2)$	
		$y = -2x + 4$	
		$2x + y - 4 = 0$	
		slope of normal $m' = -\frac{1}{m} = \frac{1}{2}$	
equation of normal is,			
$y - y_1 = m'(x - x_1)$			
$y - 0 = \frac{1}{2}(x - 2)$			
$2y = x - 2$			
$x - 2y - 2 = 0$			

	b)	Differentiate $(\sin x)^{\tan x}$ w.r.t.x	04
	Ans	Let $y = (\sin x)^{\tan x}$	
		$\log y = \tan x \log(\sin x)$	1/2
		$\frac{1}{y} \frac{dy}{dx} = \tan x \frac{1}{\sin x} \cos x + \log(\sin x) \sec^2 x$	2
		$\frac{dy}{dx} = y(\tan x \cot x + \log(\sin x) \sec^2 x)$	1
		$\frac{dy}{dx} = (\sin x)^{\tan x} (1 + \log(\sin x) \sec^2 x)$	1/2

	c)	If $Y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$, find $\frac{dy}{dx}$	04
	Ans	$y = \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}}$	



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
3.	c)	$y = \sqrt{\frac{2 \sin^2 x}{2 \cos^2 x}}$ $y = \sqrt{\tan^2 x}$ $y = \tan x$ $\frac{dy}{dx} = \sec^2 x$	1 1 1 1
	d)	<p>Evaluate: $\int \frac{\sin \sqrt{x}}{\sqrt{x}} dx$</p> <p>Ans $\int \frac{\sin \sqrt{x}}{\sqrt{x}} dx$</p> <p>Put $\sqrt{x} = t$</p> $\therefore \frac{1}{2\sqrt{x}} dx = dt$ $\therefore \frac{1}{\sqrt{x}} dx = 2dt$ $= \int \sin t (2dt)$ $= -2 \cos t + c$ $= -2 \cos \sqrt{x} + c$	04 1 ½ ½ 1½ ½
4	a)	<p>Evaluate: $\int \frac{1}{\sqrt{1-x^2} (\sin^{-1} x)^2} dx$</p> <p>Ans $\int \frac{1}{\sqrt{1-x^2} (\sin^{-1} x)^2} dx$</p> <p>Put $\sin^{-1} x = t$</p> $\therefore \frac{1}{\sqrt{1-x^2}} dx = dt$ $= \int \frac{1}{t^2} dt$ $= \int t^{-2} dt$ $= \frac{t^{-1}}{-1} + c$	12 04 1 1 ½
			1



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	a)	$= -(\sin^{-1} x)^{-1} + c$	½
	b)	Evaluate : $\int \frac{1}{5+4\cos x} dx$	04
	Ans	$\int \frac{1}{5+4\cos x} dx$ $\text{Put } \tan \frac{x}{2} = t \quad \cos x = \frac{1-t^2}{1+t^2}, dx = \frac{2dt}{1+t^2}$ $\therefore \int \frac{dx}{5+4\cos x} = \int \frac{1}{5+4\left(\frac{1-t^2}{1+t^2}\right)} \cdot \frac{2dt}{1+t^2}$ $= 2 \int \frac{1}{t^2+9} dt$ $= 2 \int \frac{1}{t^2+3^2} dt$ $= 2 \times \frac{1}{3} \tan^{-1}\left(\frac{t}{3}\right) + c$ $= \frac{2}{3} \tan^{-1}\left(\frac{\tan \frac{x}{2}}{3}\right) + c$	
c)	Evaluate: $\int \frac{x}{1+\cos 2x} dx$		
Ans	$\int \frac{x}{1+\cos 2x} dx$ $= \int \frac{x}{2\cos^2 x} dx$ $= \frac{1}{2} \int x \sec^2 x dx$ $= \frac{1}{2} \left[x \int \sec^2 x dx - \int \left(\int \sec^2 x dx \cdot \frac{d}{dx} x \right) dx \right]$ $= \frac{1}{2} \left[x \tan x - \int \tan x \cdot 1 dx \right]$ $= \frac{1}{2} \left[x \tan x - \log(\sec x) \right] + c$	½ ½ 1 1 1	



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22206

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	d)	Evaluate : $\int \frac{\sec^2 x}{(1 + \tan x)(2 + \tan x)}$	04
	Ans	$\int \frac{\sec^2 x}{(1 + \tan x)(2 + \tan x)} dx$ <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> $\int \frac{1}{(1+t)(2+t)} dt$ $\frac{1}{(1+t)(2+t)} = \frac{A}{1+t} + \frac{B}{2+t}$ $1 = A(2+t) + B(1+t)$ $\therefore \text{Put } t = -1, A = 1$ $\text{Put } t = -2, B = -1$ $\therefore \frac{1}{(1+t)(2+t)} = \frac{1}{1+t} - \frac{1}{2+t}$ $\int \frac{1}{(1+t)(2+t)} dt = \int \left(\frac{1}{1+t} - \frac{1}{2+t} \right) dt$ $= \log 1+t - \log 2+t + c$ $= \log 1 + \tan x - \log 2 + \tan x + c$ </div> <div style="width: 45%; border-left: 1px solid black; padding-left: 10px;"> <p>Put $\tan x = t$ $\therefore \sec^2 x dx = dt$</p> </div> </div>	1
			½
			½
			½
		OR	
		$\int \frac{\sec^2 x}{(1 + \tan x)(2 + \tan x)} dx$ <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> $\int \frac{1}{(1+t)(2+t)} dt$ $= \int \frac{1}{t^2 + 3t + 2} dt$ $= \int \frac{1}{t^2 + 3t + \frac{9}{4} - \frac{9}{4} + 2} dt$ $= \int \frac{1}{\left(t + \frac{3}{2}\right)^2 - \left(\frac{1}{2}\right)^2} dt$ $= \frac{1}{2} \log \left \frac{t + \frac{3}{2} - \frac{1}{2}}{t + \frac{3}{2} + \frac{1}{2}} \right + c$ $= \log \left \frac{t+1}{t+2} \right + c$ </div> <div style="width: 45%; border-left: 1px solid black; padding-left: 10px;"> <p>Put $\tan x = t$ $\therefore \sec^2 x dx = dt$</p> </div> </div>	1
			½
			½
			½
			1



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22206

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.		$= \log \left \frac{\tan x + 1}{\tan x + 2} \right + c$	½
	e)	<p>-----</p> <p>Evaluate: $\int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} dx$</p>	04
	Ans	$I = \int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} dx \text{-----(1)}$ $= \int_0^{\pi/2} \frac{\sqrt[3]{\sin\left(\frac{\pi}{2} - x\right)}}{\sqrt[3]{\cos\left(\frac{\pi}{2} - x\right)} + \sqrt[3]{\sin\left(\frac{\pi}{2} - x\right)}} dx$	1
		$I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} dx \text{-----(2)}$	½
		<p>Add (1) and (2)</p> $\therefore 2I = \int_0^{\pi/2} \frac{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} dx$ $= \int_0^{\pi/2} 1 \cdot dx$ $= [x]_0^{\pi/2}$ $2I = \frac{\pi}{2}$ $\therefore I = \frac{\pi}{4}$	½
		<p>OR</p> <div style="display: flex; align-items: center;"> <div style="flex: 1;"> $I = \int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} dx$ $\therefore I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} dx$ $\therefore 2I = \int_0^{\pi/2} \frac{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} dx$ $= \int_0^{\pi/2} 1 \cdot dx$ </div> <div style="border-left: 1px solid black; padding-left: 10px; margin-left: 10px;"> <p>Replace $x \rightarrow \frac{\pi}{2} - x$</p> <p>$\therefore \sin x \rightarrow \cos x$</p> <p>& $\cos x \rightarrow \sin x$</p> </div> </div>	1
		$= \int_0^{\pi/2} 1 \cdot dx$	½



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
4.	e)	$= [x]_0^{\pi/2}$ $2I = \frac{\pi}{2}$ $\therefore I = \frac{\pi}{4}$	1
5	a)	<p>Attempt any TWO of the following:</p> <p>Find the area of the region bounded by the parabola $y = 4x - x^2$ and the x-axis.</p>	12
		<p>Ans $y = 4x - x^2$</p> <p>put $y = 0$,</p> $4x - x^2 = 0$ $x = 0, x = 4$	06
		$\text{Area} = \int_a^b y dx$ $= \int_0^4 (4x - x^2) dx$	1
		$= 4 \left[\frac{x^2}{2} \right]_0^4 - \left[\frac{x^3}{3} \right]_0^4$ <p style="text-align: center;">OR $= \left[4 \frac{x^2}{2} - \frac{x^3}{3} \right]_0^4$</p>	2
		$= 4 \left[\frac{4^2}{2} - \frac{0^2}{2} \right] - \left[\frac{4^3}{3} - \frac{0^3}{3} \right]$ <p style="text-align: center;">OR $= \left[\left(2(4)^2 - \frac{4^3}{3} \right) - 0 \right]$</p> $= \frac{32}{3} = 10.667$	1
b)	<p>Attempt the following:</p> <p>(i) Form the D.E. by eliminating the arbitrary constants if $y = A \cos 3x + B \sin 3x$</p>	06	
Ans	<p>$y = A \cos 3x + B \sin 3x$</p> $\therefore \frac{dy}{dx} = -3A \sin 3x + 3B \cos 3x$ $\therefore \frac{d^2y}{dx^2} = -9A \cos 3x - 9B \sin 3x$ $\therefore \frac{d^2y}{dx^2} = -9(A \cos 3x + B \sin 3x)$ $\frac{d^2y}{dx^2} = -9y$ $\frac{d^2y}{dx^2} + 9y = 0$	03	
			1
			1
			½
			½



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
5.	b)(ii)	Solve : $x(1+y^2)dx + y(1+x^2)dy = 0$	03
	Ans	$x(1+y^2)dx + y(1+x^2)dy = 0$ $\therefore \frac{x}{1+x^2} dx = -\frac{y}{1+y^2} dy$ $\therefore \int \frac{x}{1+x^2} dx = -\int \frac{y}{1+y^2} dy$ $\therefore \frac{1}{2} \log(1+x^2) = -\frac{1}{2} \log(1+y^2) + c$ $\therefore \log(1+x^2) = -\log(1+y^2) + C$ <p>-----</p>	1 1 1
	(c)	A particle starting with velocity 6m/sec has an acceleration $(1-t^2)$ m/sec ² , when does it first come to rest? How far has it then travelled?	06
	Ans	<p>Acceleration = $\frac{dv}{dt} = 1-t^2$</p> $\therefore dv = (1-t^2) dt$ $\therefore \int dv = \int (1-t^2) dt$ $\therefore v = t - \frac{t^3}{3} + c$ <p>given $v = 6$ and initially $t = 0$</p> $\therefore c = 6$ $\therefore v = t - \frac{t^3}{3} + 6$ <p>The particle comes to rest when $v = 0$</p> $\therefore t - \frac{t^3}{3} + 6 = 0$ $\therefore t^3 - 3t - 18 = 0$ $\therefore t = 3$ $\therefore v = \frac{dx}{dt}$ $\therefore \frac{dx}{dt} = t - \frac{t^3}{3} + 6$ $\therefore dx = \left(t - \frac{t^3}{3} + 6 \right) dt$ $\therefore \int dx = \int \left(t - \frac{t^3}{3} + 6 \right) dt$	1 ½ ½ 1 ½



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22206

Q. No.	Sub Q. N.	Answer	Marking Scheme
5.	c)	$\therefore x = \frac{t^2}{2} - \frac{t^4}{12} + 6t + c_1$	1
		$\therefore \text{initially } x = 0, t = 0$ $c_1 = 0$ $\therefore x = \frac{t^2}{2} - \frac{t^4}{12} + 6t$ <p>put $t = 3$</p> $\therefore x = \frac{(3)^2}{2} - \frac{(3)^4}{12} + 6(3)$ $\therefore x = 15.75$	1/2
6		Attempt any TWO of the following:	12
	a)	Attempt the following:	06
	i)	A person fires 10 shots at target. The probability that any shot will hit the target $3/5$. Find the probability that the target is hit exactly 5 times.	03
	Ans	$n = 10, p = \frac{3}{5}$ $q = 1 - p = 1 - \frac{3}{5} = \frac{2}{5}$ $r = 5$ $p(r) = {}^n C_r (p)^r (q)^{n-r}$ $p(5) = {}^{10} C_5 \left(\frac{3}{5}\right)^5 \left(\frac{2}{5}\right)^{10-5}$ $= 0.2007$	2 1
ii)	If 20% of the bolt produce by a machine are defective .Find the Probability that out of 4 bolts drawn , (1) one is defective (2) at the most two are defective.	03	
Ans	<p>Given $p = 20\% = \frac{20}{100} = 0.2, n = 4$ and $q = 1 - p = 0.8$</p> $p(r) = {}^n C_r p^r q^{n-r}$		



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22206**

Q. No.	Sub Q. N.	Answer	Marking Scheme
6.	a)(ii)	(1) p (one is defective) $= p(1) = 4C_1 (0.2)^1 (0.8)^{4-1}$ $= 0.4096$	1 ½
		(2) p (at the most two are defective.) $= p(0) + p(1) + p(2)$ $= 4C_0 (0.2)^0 (0.8)^{4-0} + 4C_1 (0.2)^1 (0.8)^{4-1} + 4C_2 (0.2)^2 (0.8)^{4-2}$ $= 0.9728$	1 ½

	b)	A company manufacture electric motors. The probability that an electric motor is defective is 0.01. What is the probability that a sample of 300 electric motors will contain exactly 5 defective motors? (Given: $e^{-3} = 0.0498$)	06
	Ans	$p = 0.01, n = 300, r = 5$ $\therefore m = np = 0.01 \times 300 = 3$ $p(r) = \frac{e^{-m} \cdot (m)^r}{r!}$ $p(5) = \frac{e^{-3} \cdot (3)^5}{5!}$ $p(5) = \frac{(0.0498) \cdot (3)^5}{5!}$ $= 0.1008$	2 2 1 1
	c)	In a sample of 1000 cases the mean of certain test is 14 and standard deviation is 2.5. Assuming the distribution to be normal, find (1) how many students score above 18? (2) how many students score between 12 and 15? [Given: $A(0.4) = 0.1554, A(0.8) = 0.2881, A(1.6) = 0.4452$]	06
	Ans	Given $\bar{x} = 14, \sigma = 2.5, N = 1000$ (1) $z = \frac{x - \bar{x}}{\sigma} = \frac{18 - 14}{2.5} = 1.6$ $\therefore p(\text{score above } 18) = A(\text{greater than } 1.6)$ $= 0.5 - A(1.6)$ $= 0.5 - 0.4452 = 0.0548$ $\therefore \text{No. of students} = N \cdot p$ $= 1000 \times 0.0548 = 54.8 \text{ i.e., } 55$	1 1 1

